Goal: End the Epidemic

Develop coordinated clinical research plan

Goal: End the Epidemic Develop coordinated clinical research plan

Stop new infections

- Protect uninfected persons
- Reduce infectiousness of infected persons
- Keep infected persons healthy
- Continue to pursue innovative translational research

Issues To Be Considered

Continuum of Approaches

- Applicability

 - Timelines
- Individual RCT ← community RCT
- Domestic ← international 'agenda'
- "Long term clinical outcome ↔ "surrogate markers"

Issues To Be Considered

Continuum of Approaches

- Routes of exposure
 - Vaginal, rectal, perinatal (peripartum, breastfeeding), parenteral, oral
- Ages
 - Adults, adolescents, newborns, perinatal
- Focus?

- Protect uninfected
 - Vaccines
 - Microbicides
 - Therapies/vaccines
 - Coinfections, particularly genital ulcerative infections
 - Behavior
 - Barrier methods
- Reduce infectiousness of infected
 - Reduce viral load and/or shedding
 - ART, co-infection treatment, vaccines?
 - Modify transmission events
 - Behavior, barrier methods

- HIV Vaccines issues
 - Multi-country trials
 - Clade issue
 - Background issue (HLA/genetic, other)
 - How many countries/populations?

- Vaccines
 - 2-3 efficacy trials; impact on infection, disease progression and secondary transmission
 - RCT and also community based?
 - Selection and improvement
 - Selection of most promising candidates
 - Approaches to compare candidates, e.g. pox vectors
 - Continued improvement in vaccine design
 - Antibody !!!

- Passive transfer in MTCT
 - Selection of best possible combinations for "proof of concept" in MTCT setting

- Microbicides issues
 - Adherence and selection of control arm(s)
- Science
 - 1-2 phase III trials
 - RCT
 - Placebo vs. condom-only arm as comparison
 - Selection of most promising products
 - New microbicide design

- Behavior (individual, community)
 - Impact VCT on transmission in RLDC
- Barriers (male vs. female)
- Treatments (ART, Co-infections)
- Vaccines (HPV?)

Keep Infected Persons Healthy

- When to start and with what; when to switch
 - Early initiation or deferred therapy
 - Adherence, cost-benefit, drug conservation
- Resistance and salvage
 - Sequencing regimens to retard resistance
 - New approaches for highly treated new RX
- Immune preservation or restoration
 - Immune modulation to augment lymphocyte number or function
 - "Therapeutic" vaccines

Keep Infected Persons Healthy

- Co-infections/Concomitant Meds
 - Tuberculosis
 - Hepatitis B, C and GBVc
 - Malaria
 - Natural products
- Reduce or control complications
 - Managing CV, GI and neuro signs and sx
 - Developing new drug regimens
- Simplify diagnosis, monitoring, delivery

- Common trials single trial jointly planned and conducted
 - AVEG/HPTN phase II trial
- Simultaneous trials
 - HPTN052, AACTG 5175, AACTG 5190
- Trials conducted in the same setting
- Data sharing

- Impact of prevention and ART on transmission
- Rx of HIV+ women in RLDC and impact on MTCT
 - Research relevant to family members
- Rx and prevention vaccine research
 - Common correlates?

- Studies of acute infection; viral dynamics; reservoirs
 - Vaccines, Rx, microbicides
- Cross-cutting lab research
 - Diagnosis, endpoint monitoring (viral load, CD4), immune assays, other?
- Cross-cutting background issues
 - Nutrition, traditional therapies, endemic infections

- Impact of prevention and ART on transmission
- Rx of HIV+ women in RLDC; impact on MTCT
- Rx and prevention vaccine research
- Studies of acute infection
- Cross-cutting lab research
- Cross-cutting background issues
- > YOUR IDEAS? PRIORITIES?